



Some remarks on $\Delta^m(I_\lambda)$ –summability on neutrosophic normed spaces

Archana Sharma¹, Sajid Murtaza², Vijay Kumar^{3,*}

^{1,2,3} Department of Mathematics, Chandigarh University, Gharuan Mohali (Punjab), India

Emails: dr.archanasharma1022@gmail.com; sajidsulimani8@gmail.com; kaushikvjy@gmail.com

Abstract

In the present paper, we use the difference operator Δ^m and λ – summability to define some new summability concepts on neutrosophic normed spaces. We also introduce concepts of generalized limit point, and cluster point and obtain some relationships among these notions. Finally, we define generalized Cauchy sequences on these spaces and present a characterization of a new summability method that preserves linear operators on neutrosophic normed spaces.

Keywords: Neutrosophic normed spaces; lacunary convergence; and I -convergence.

1. Introduction

Statistical convergence was defined by Fast [9] and further studied in [5], [10], [11], [24] and [25]. “A sequence (x_k) of numbers is said to be statistical convergence to a number L if for each $\varepsilon > 0$, $\lim_n \frac{1}{n} |\{k \leq n: |x_k - L| \geq \varepsilon\}| = 0$ or $\delta(K_\varepsilon) = 0$, where K_ε by $\{k \leq n: |x_k - L| \geq \varepsilon\} \subseteq \mathbb{N}$ ”.

In [20], the (V, λ) – summability is defined as follows: “For any non-decreasing sequence of positive numbers $\lambda = (\lambda_n)$ with $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$ and $\lambda_{n+1} \leq \lambda_n + 1$, $\lambda_1 = 1$, the generalized de la Valée-Pousin mean is defined by

$$t_n(x) = \frac{1}{\lambda_n} \sum_{k \in I_n} x_k \quad \text{where } I_n = [n - \lambda_n + 1, n].$$

A sequence $x = (x_k)$ is said to be λ – summable to a number L if $t_n(x) = \frac{1}{\lambda_n} \sum_{k \in I_n} x_k \rightarrow L$ as $n \rightarrow \infty$ ”.

Statistical convergence is further generalized in [16], called I – convergence. Later, the idea is developed in [4], [6], [7], [8], [12], [15], [17], [18], [19] and [21]. For basic information on I – convergence, we recommend the paper [16].

Fuzzy sets were invented by Zadeh [28] and generalized by Atanassov [1], called intuitionistic fuzzy sets. Over the years, many applications of these sets can be found in [6], [12], [21], [22] and [23]. Recently, Smarandache [27] introduced a generalization of an intuitionistic fuzzy set, called a neutrosophic set. For some recent works on these sets, we refer [2], [3], [13] and [14]. We aim in this paper, to define $\Delta^m(\lambda)$ – convergence and $\Delta^m(I_\lambda)$ – convergence on neutrosophic normed spaces as a generalization of ideal convergence. Later, we introduce the concepts of $\Delta^m(I_\lambda)$ – limit point,

$\Delta^m(I_\lambda)$ –cluster point and obtain some relationships among these. We also define $\Delta^m(\lambda)$ – Cauchy, $\Delta^m(I_\lambda)$ –Cauchy sequences on these spaces and study their relations. Finally, we present a characterization for $\Delta^m(I_\lambda)$ –convergence preserving linear operators on neutrosophic normed spaces

2. Background and Preliminary

In this section, we review and give some definitions and results which form the basis for the present study.

Definition 2.1[16] “Let F be a vector space, $\mathcal{N} = \{(\vartheta, \mathcal{G}(\vartheta), \mathcal{B}(\vartheta), \mathcal{Y}(\vartheta)) : \vartheta \in F\}$ be a normed space such that $\mathcal{N} : F \times \mathbb{R}^+ \rightarrow [0, 1]$ and \circ, \bullet respectively are continuous t – norm and continuous t – conorm. Then a four tuple $V = (F, \mathcal{N}, \circ, \bullet)$ is called a neutrosophic normed space (NNS) if the following conditions are satisfied.

For every $u, v \in F$ and $\lambda, \mu > 0$ and for every $\sigma \neq 0$ we have

- (i) $0 \leq \mathcal{G}(u, \lambda) \leq 1, 0 \leq \mathcal{B}(u, \lambda) \leq 1, 0 \leq \mathcal{Y}(u, \lambda) \leq 1$ for every $\lambda \in \mathbb{R}^+$;
- (ii) $\mathcal{G}(u, \lambda) + \mathcal{B}(u, \lambda) + \mathcal{Y}(u, \lambda) \leq 3$ for $\lambda \in \mathbb{R}^+$;
- (iii) $\mathcal{G}(u, \lambda) = 1$ (for $\lambda > 0$) if and only if $u = 0$;
- (iv) $\mathcal{G}(\sigma u, \lambda) = \mathcal{G}\left(u, \frac{\lambda}{|\sigma|}\right)$;
- (v) $\mathcal{G}(u, \mu) \circ \mathcal{G}(v, \lambda) \leq \mathcal{G}(u + v, \lambda + \mu)$;
- (vi) $\mathcal{G}(u, \cdot)$ is a continuous non-decreasing function;
- (vii) $\lim_{\lambda \rightarrow \infty} \mathcal{G}(u, \lambda) = 1$;
- (viii) $\mathcal{B}(u, \lambda) = 0$ (for $\lambda > 0$) if and only if $u = 0$;
- (ix) $\mathcal{B}(\sigma u, \lambda) = \mathcal{B}\left(u, \frac{\lambda}{|\sigma|}\right)$;
- (x) $\mathcal{B}(u, \mu) \bullet \mathcal{B}(v, \lambda) \geq \mathcal{B}(u + v, \lambda + \mu)$;
- (xi) $\mathcal{B}(u, \cdot)$ is a continuous non-decreasing function;
- (xii) $\lim_{\lambda \rightarrow \infty} \mathcal{B}(u, \lambda) = 0$;
- (xiii) $\mathcal{Y}(u, \lambda) = 0$ (for $\lambda > 0$) if and only if $u = 0$;
- (xiv) $\mathcal{Y}(\sigma u, \lambda) = \mathcal{Y}\left(u, \frac{\lambda}{|\sigma|}\right)$;
- (xv) $\mathcal{Y}(u, \mu) \bullet \mathcal{Y}(v, \lambda) \geq \mathcal{Y}(u + v, \lambda + \mu)$;
- (xvi) $\mathcal{Y}(u, \cdot)$ is a continuous non-decreasing function;
- (xvii) $\lim_{\lambda \rightarrow \infty} \mathcal{Y}(u, \lambda) = 0$ and
- (xviii) If $\lambda \leq 0$, then $\mathcal{G}(u, \lambda) = 0, \mathcal{B}(u, \lambda) = 1$ and $\mathcal{Y}(u, \lambda) = 1$.

Here, $\mathcal{N}(\mathcal{G}, \mathcal{B}, \mathcal{Y})$ is called the neutrosophic norm”.

Some examples of neutrosophic normed spaces can be found in [16].

“A sequence (a_k) in Neutrosophic Normed Spaces V is said to *convergent* if, for each $\varepsilon > 0$ and $\lambda > 0$, there exists a positive integer m and $\mathcal{L} \in F$ such that

$$\mathcal{G}(a_k - \mathcal{L}, \lambda) > 1 - \varepsilon, \mathcal{B}(a_k - \mathcal{L}, \lambda) < \varepsilon \text{ and } \mathcal{Y}(a_k - \mathcal{L}, \lambda) < \varepsilon \text{ for all } k \geq m.$$

This is equivalent to say

$$\lim_{k \rightarrow \infty} \mathcal{G}(a_k - \mathcal{L}, \lambda) = 1, \lim_{k \rightarrow \infty} \mathcal{B}(a_k - \mathcal{L}, \lambda) = 0 \text{ and } \lim_{k \rightarrow \infty} \mathcal{Y}(a_k - \mathcal{L}, \lambda) = 0.$$

and we write in this case $\mathcal{N} - \lim_{k \rightarrow \infty} a_k = \mathcal{L}$.”

“The sequence (a_k) is said to be *Cauchy* if, for each $\varepsilon > 0$ and $\lambda > 0$, there exists a positive integer m and such that

$\mathcal{G}(a_k - a_n, \lambda) > 1 - \varepsilon, \mathcal{B}(a_k - a_n, \lambda) < \varepsilon$ and $\mathcal{Y}(a_k - a_n, \lambda) < \varepsilon$ for all $k, n \geq m$."

For basic terminology on neutrosophic normed space, we refer to [14].

Let w denotes the set of all sequences in the neutrosophic normed space $V = (F, \mathcal{N}, \circ, \bullet)$. Define $\Delta^m: w \rightarrow w$ by

$$\begin{aligned} \Delta^0 a_k &= a_k; \\ \Delta^1 a_k &= a_k - a_{k+1}; \\ \Delta^m a_k &= \Delta^{m-1} (\Delta a_k) = \Delta^{m-1} (a_k - a_{k+1}) \quad m \geq 2 \quad \text{and for all } k \in \mathbb{N}. \end{aligned}$$

Throughout the work V be an NNS ; $\lambda = (\lambda_n)$ is a sequence as described above and $I \subseteq \wp(\mathbb{N})$ denotes an admissible ideal.

3. $\Delta^m(\lambda)$ –Convergence

Definition 3.1 Let $0 < \varepsilon < 1$ and $\mu > 0$. A sequence $x = (x_k)$ in V is said to be $\Delta^m(\lambda)$ –convergent in neutrosophic norm \mathcal{N} if $\exists, \mathcal{L} \in F$ and an $n_0 \in \mathbb{N}$ satisfying

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) &> 1 - \varepsilon, \\ \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) &< \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{for } n \geq n_0, \end{aligned}$$

and we write $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}$.

Theorem 3.1 If $x = (x_k)$ be a $\Delta^m(\lambda)$ –convergent with $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}$, then, \mathcal{L} is unique.

Proof Assume, there exists \mathcal{L}_1 and \mathcal{L}_2 in V such that $\mathcal{L}_1 \neq \mathcal{L}_2$ and $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}_1, \mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}_2$. let $\varepsilon > 0$ and $\mu > 0$. Choose $\vartheta > 0$ such that

$$(1 - \varepsilon) \circ (1 - \varepsilon) > 1 - \vartheta \quad \text{and} \quad \varepsilon \bullet \varepsilon < \vartheta \tag{3.1}$$

Since, $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}_1, \mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}_2$ so for every $\varepsilon > 0$ and $\mu > 0 \exists n_1$ and n_2 in \mathbb{N} with

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_1, \mu) &> 1 - \varepsilon, \\ \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_1, \mu) &< \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_1, \mu) < \varepsilon \quad \text{for } n \geq n_1; \end{aligned}$$

and

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(x \Delta^m_k - \mathcal{L}_2, \mu) &> 1 - \varepsilon, \\ \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_2, \mu) &< \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_2, \mu) < \varepsilon \quad \text{for } n \geq n_2. \end{aligned}$$

Case (i) $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_1, \mu) > 1 - \varepsilon$ and $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_2, \mu) > 1 - \varepsilon;$

Case (ii) $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_1, \mu) < \varepsilon$ and $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_2, \mu) < \varepsilon$ and

Case (iii) $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_1, \mu) < \varepsilon$ and $\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_2, \mu) < \varepsilon$.

Case (i) Let $n_0 = \max\{n_1, n_2\}$, then for $n \geq n_0$ we can find $p \in \mathbb{N}$ satisfying

$$\mathcal{G}\left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2}\right) > \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}\left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2}\right) > 1 - \varepsilon \quad \text{and}$$

$$\mathcal{G}\left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2}\right) > \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}\left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2}\right) > 1 - \varepsilon.$$

Now,

$$\mathcal{G}(\mathcal{L}_1 - \mathcal{L}_2, \mu) \geq \mathcal{G}\left(\Delta^m a_p - \mathcal{L}_1, \frac{\mu}{2}\right) \circ \mathcal{G}\left(\Delta^m a_p - \mathcal{L}_2, \frac{\mu}{2}\right) > (1 - \varepsilon) \circ (1 - \varepsilon) > 1 - \vartheta \tag{3.2}$$

Since ϑ is arbitrary and (3.2) holds for every $\mu > 0$, it follows that $\mathcal{G}(\mathcal{L}_1 - \mathcal{L}_2, \mu) = 1$, and therefore $\mathcal{L}_1 = \mathcal{L}_2$.

Case (ii) As in case (i) for $n \geq n_0$ there exists $p \in \mathbb{N}$ such that

$$\mathcal{B}\left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2}\right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}\left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2}\right) < \varepsilon \quad \text{and}$$

$$\mathcal{B}\left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2}\right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}\left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2}\right) < \varepsilon.$$

Now,

$$\mathcal{B}(\mathcal{L}_1 - \mathcal{L}_2, \mu) < \mathcal{B}\left(\Delta^m a_p - \mathcal{L}_1, \frac{\mu}{2}\right) \circ \mathcal{B}\left(\Delta^m a_p - \mathcal{L}_2, \frac{\mu}{2}\right) < \varepsilon \bullet \varepsilon < \vartheta.$$

Since ϑ is arbitrary and above inequality holds for every $\mu > 0$, it follows that $\mathcal{B}(\mathcal{L}_1 - \mathcal{L}_2, \mu) = 0$, and therefore $\mathcal{L}_1 = \mathcal{L}_2$.

Case (iii) follows similarly to case (ii) ■

Theorem 3.2 Let $x = (x_k)$, $y = (y_k)$ be two sequences in V such that $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}_1$ and $\mathcal{N}_\lambda - \lim_k \Delta^m y_k = \mathcal{L}_2$, then

- (i) $\mathcal{N}_\lambda - \lim_k (\Delta^m (x_k + y_k)) = \mathcal{L}_1 + \mathcal{L}_2$.
- (ii) $\mathcal{N}_\lambda - \lim_k (\beta (\Delta^m x_k)) = \beta \mathcal{L}_1$ for $\beta \neq 0$

Proof. excluded. ■

Theorem 3.3 For any $x = (x_k)$ with $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}$, \exists a subsequence (x_{k_j}) of (x_k) with $\mathcal{N} - \lim_j \Delta^m x_{k_j} = \mathcal{L}$.

Proof Suppose $\mathcal{N}_\lambda - \lim_k \Delta^m x_k = \mathcal{L}$ holds. For each $\varepsilon > 0$ and $\mu > 0$, $\exists n_0 \in \mathbb{N}$ satisfying

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon,$$

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{for } n \geq n_0.$$

Now for $n \geq n_0$, we can select a $k_j \in I_n$ such that

$$\mathcal{G}\left(\Delta^m x_{k_j} - \mathcal{L}, \mu\right) > \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon;$$

$$\mathcal{B}(\Delta^m x_{k_j} - \mathcal{L}, \mu) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{and}$$

$$\mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon.$$

Thus, we have a subsequence (x_{k_j}) of (x_k) such that $\mathcal{N} - \lim_j \Delta^m x_{k_j} = \mathcal{L}$. ■

4. $\Delta^m(I_\lambda)$ – Convergence

Definition 4.1, Let $0 < \varepsilon < 1$ and $\mu > 0$. A sequence $x = (x_k)$ in V is said to be $\Delta^m(I_\lambda)$ – convergent with respect to \mathcal{N} if $\exists, \mathcal{L} \in F$ satisfying

$$A(\varepsilon, \mu) = \left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \in I.$$

We denote it by $\Delta^m(I_\lambda^{\mathcal{N}}) - \lim_k x_k = \mathcal{L}$.

Example 4.1 Let $(F, \|\cdot\|)$ be a normed space. Define t – norm \circ and t – conorm \bullet as follows. For $u, v \in [0, 1]$, $u \circ v = uv$ and $u \bullet v = \min\{u + v, 1\}$. For $\mu > 0$ and $\mu > \|u\|$, define

$$\mathcal{G}(u, \lambda) = \frac{\mu}{\mu + \|u\|}, \quad \mathcal{B}(u, \lambda) = \frac{\|u\|}{\mu + \|u\|} \quad \text{and} \quad \mathcal{Y}(u, \lambda) = \frac{\|u\|}{\mu}$$

Then, $(F, \mathcal{N}, \circ, \bullet)$ is a neutrosophic normed space.

Let, $m = 0$, $\lambda = (\lambda_n) = (n)$ and chose $I = \{S \subseteq \mathbb{N} : \delta(S) = 0\}$. Construct a sequence $x = (x_k)$ by

$$x_k = \begin{cases} 1 & \text{if } k = j^2 \text{ where } j \in \mathbb{N}; \\ 0, & \text{otherwise;} \end{cases}$$

Then for $\varepsilon > 0$ and $\mu > 0$, the set

$$A(\varepsilon, \mu) = \left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k, \mu) \geq \varepsilon \right\}$$

is contained in the set of squares whose natural density is zero. So, we have $\delta(A(\varepsilon, \mu)) = 0$ and therefore $\delta(A(\varepsilon, \mu)) \in I$. ■

Lemma 4.1 For $\varepsilon > 0$ and $\mu > 0$, the following are equivalents.

- (i) $\Delta^m(I_\lambda^{\mathcal{N}}) - \lim_k x_k = \mathcal{L}$;
- (ii) $\left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \right\} \in I$; $\left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \in I$ and $\left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \in I$.

- (iii) $\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon, \right.$
 $\left. \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \right\}$
 $\in \mathcal{F}(I)$
- (iv) $\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon \right\}$
 $\in \mathcal{F}(I); \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \right\} \in \mathcal{F}(I) \text{ and}$
 $\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \right\} \in \mathcal{F}(I)$
- (v) $I_\lambda - \lim_k \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) = 1; \quad I_\lambda - \lim_k \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) = 0 \quad \text{and } I_\lambda$
 $- \lim_k \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) = 0.$

Proof. Omitted ■

Next Theorem provided the uniqueness of $I_N^\theta - \lim_k x_k$ provided it exists.

Theorem 4.1 If $x = (x_k)$ is a sequence in V such that $\Delta^m(I_N^\lambda) - \lim_k x_k = \mathcal{L}_1$ and $\Delta^m(I_N^\lambda) - \lim_k x_k = \mathcal{L}_2$, then $\mathcal{L}_1 = \mathcal{L}_2$.

Proof: Suppose that $\mathcal{L}_1 \neq \mathcal{L}_2$ and let $\varepsilon > 0$. Choose $\gamma > 0$ such that

$$(1 - \varepsilon) \circ (1 - \varepsilon) > 1 - \gamma \quad \text{and} \quad \varepsilon \bullet \varepsilon < \gamma \tag{4.1}$$

For $\mu > 0$, define.

$$K_{\mathcal{G}_1}(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) \leq 1 - \varepsilon \right\},$$

$$K_{\mathcal{G}_2}(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) \leq 1 - \varepsilon \right\};$$

$$K_{\mathcal{B}_1}(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) \geq \varepsilon \right\}, \quad K_{\mathcal{B}_2}(\varepsilon, \mu)$$

$$= \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) \geq \varepsilon \right\};$$

$$K_{\mathcal{Y}_1}(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) \geq \varepsilon \right\} \text{ and } K_{\mathcal{Y}_2}(\varepsilon, \mu)$$

$$= \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) \geq \varepsilon \right\}.$$

Since $\Delta^m(I_N^\lambda) - \lim_k x_k = \mathcal{L}_1$ and $\Delta^m(I_N^\lambda) - \lim_k x_k = \mathcal{L}_2$ so by Lemma 3.1, we respectively have the sets $K_{G_1}(\varepsilon, \mu)$; $K_{B_1}(\varepsilon, \mu)$; $K_{y_1}(\varepsilon, \mu)$ and $K_{G_2}(\varepsilon, \mu)$; $K_{B_2}(\varepsilon, \mu)$ and $K_{y_2}(\varepsilon, \mu)$ belongs to I . Define a set $K_N(\varepsilon, \mu)$ by

$$K_N(\varepsilon, \mu) = \left\{ \left\{ \left\{ K_{G_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{G_2}(\varepsilon, \mu) \right\} \right\} \cap \left\{ \left\{ K_{B_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{B_2}(\varepsilon, \mu) \right\} \right\} \right. \\ \left. \cap \left\{ \left\{ K_{y_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{y_2}(\varepsilon, \mu) \right\} \right\} \right\};$$

then $K_N(\varepsilon, \mu) \in I$ and therefore $\emptyset \neq K_N^c(\varepsilon, \mu) \in \mathcal{F}(I)$. Let $n \in K_N^c(\varepsilon, \mu)$, then

- (i) $n \in \mathbb{N} - \left\{ \left\{ K_{G_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{G_2}(\varepsilon, \mu) \right\} \right\}$
- (ii) $n \in \mathbb{N} - \left\{ \left\{ K_{B_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{B_2}(\varepsilon, \mu) \right\} \right\}$
- (iii) $n \in \mathbb{N} - \left\{ \left\{ K_{y_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{y_2}(\varepsilon, \mu) \right\} \right\}$.

Suppose that (i) holds, then $n \notin \left\{ K_{G_1}(\varepsilon, \mu) \right\} \cup \left\{ K_{G_2}(\varepsilon, \mu) \right\}$ which gives $n \notin K_{G_1}(\varepsilon, \mu)$ and $n \notin K_{G_2}(\varepsilon, \mu)$. This implies that

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) > 1 - \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) > 1 - \varepsilon \tag{4.2}$$

Clearly, we will get a $p \in \mathbb{N}$ such that

$$\mathcal{G} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) > \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) > 1 - \varepsilon \quad \text{and} \\ \mathcal{G} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) > \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) > 1 - \varepsilon.$$

Now,

$$\mathcal{G}(\mathcal{L}_1 - \mathcal{L}_2, \mu) \geq \mathcal{G} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) \circ \mathcal{G} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) > (1 - \varepsilon) \circ (1 - \varepsilon) > 1 - \gamma \tag{4.3}$$

(using (4.1) and (4.2))

Since γ is arbitrary and (4.3) holds for every $\mu > 0$, it follows that $\mathcal{G}(\mathcal{L}_1 - \mathcal{L}_2, \mu) = 1$, and therefore $\mathcal{L}_1 = \mathcal{L}_2$.

We now assume (ii) holds, then $n \notin K_{B_1}(\varepsilon, \mu)$ and $n \notin K_{B_2}(\varepsilon, \mu)$, and therefore we have

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) < \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon \tag{4.4}$$

Using the same technique as above, we get $p \in \mathbb{N}$ such that

$$\mathcal{B} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) < \varepsilon \quad \text{and} \\ \mathcal{B} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon.$$

Now,

$$\mathcal{B}(\mathcal{L}_1 - \mathcal{L}_2, \lambda) \leq \mathcal{B} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) \bullet \mathcal{B} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon \bullet \varepsilon < \gamma \tag{4.5}$$

(using (4.1) and (4.4))

As γ is arbitrary and (4.5) holds for every $\mu > 0$, we must have $\mathcal{B}(\mathcal{L}_1 - \mathcal{L}_2, \mu) = 0$, which gives immediately $\mathcal{L}_1 = \mathcal{L}_2$.

Finally, assume that (iii) holds. It follows that $n \notin K_{y_1}(\mu, \lambda)$, and $n \notin K_{y_2}(\mu, \lambda)$ and therefore we have

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) < \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon$$

As above, we get $p \in \mathbb{N}$ such that

$$\mathcal{Y} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_1, \frac{\mu}{2} \right) < \varepsilon \quad \text{and}$$

$$\mathcal{Y} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) < \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y} \left(\Delta^m x_k - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon.$$

Now,

$$\mathcal{Y}(\mathcal{L}_1 - \mathcal{L}_2, \mu) \leq \mathcal{Y} \left(\Delta^m x_p - \mathcal{L}_1, \frac{\mu}{2} \right) \cdot \mathcal{Y} \left(\Delta^m x_p - \mathcal{L}_2, \frac{\mu}{2} \right) < \varepsilon \cdot \varepsilon < \gamma \tag{4.6}$$

Since γ is arbitrary and (4.6) holds for every $\mu > 0$, we must have $\mathcal{Y}(\mathcal{L}_1 - \mathcal{L}_2, \mu) = 0$ and therefore we have $\mathcal{L}_1 = \mathcal{L}_2$. ■

Theorem 4.2 If $x = (x_k), y = (y_k)$ be two sequences in V such that $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k x_k = \mathcal{L}_1$ and $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k y_k = \mathcal{L}_2$, then

- (i) $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k (x_k + y_k) = \mathcal{L}_1 + \mathcal{L}_2$
- (ii) $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k (\beta x_k) = \beta \mathcal{L}_1$ for $\beta \neq 0$

Proof. Omitted. ■

Next Theorem provide the relation between $\Delta^m(\mathcal{N}_\lambda) - convergence$ and $\Delta^m(I_{\mathcal{N}}^\lambda) - convergence$.

Theorem 4.3 For $x = (x_k)$ if $\Delta^m(\mathcal{N}_\lambda) - \lim_k x_k = \mathcal{L}$, then, $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k x_k = \mathcal{L}$.

Proof Suppose $\Delta^m(\mathcal{N}_\lambda) - \lim_k x_k = \mathcal{L}$ holds. Then for $\varepsilon > 0$ and $\mu > 0, \exists n_0 \in \mathbb{N}$ s.t

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{and} \quad \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \quad \text{for } n \geq n_0.$$

Let,

$$A(\varepsilon, \mu) = \left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \right\},$$

then it is clear that $A(\varepsilon, \mu) \subseteq \{1, 2, 3, \dots, n_0 - 1\}$ so in I . This implies $\Delta^m(I_{\mathcal{N}}^\lambda) - \lim_k x_k = \mathcal{L}$. ■

Theorem 4.4 If $x = (x_k)$ be $\Delta^m(\lambda) - convergent$ in V and $y = (y_k)$ be another sequence in V such that $\{n \in \mathbb{N} : \Delta^m x_k \neq \Delta^m y_k \text{ for some } k \in I_n\} \in I$, then (y_k) is $\Delta^m(I_\lambda) - convergent$ to the same limit.

Proof Let $\Delta^m(\mathcal{N}_\lambda) - \lim_k x_k = \mathcal{L}$. For each $\varepsilon > 0$ and $\mu > 0, \exists n_0 \in \mathbb{N}$ s.t

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon ,$$

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \text{ for } n \geq n_0 .$$

Thus

$$A(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \subseteq \{1, 2, 3, \dots, n_0 - 1\} \in I .$$

This implies that

$$A^c(\varepsilon, \mu) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}, \mu) < \varepsilon \right\} \notin I .$$

Now, for $0 < \varepsilon < 1$ and $\mu > 0$ we have

$$\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m y_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m y_k - \mathcal{L}, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m y_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \subseteq \{n \in \mathbb{N}: \Delta^m x_k \neq \Delta^m y_k \text{ for some } k \in I_n\} \cup A(\varepsilon, \mu) .$$

Since, $A(\varepsilon, \mu)$ and $\{n \in \mathbb{N}: \Delta^m x_k \neq \Delta^m y_k \text{ for some } k \in I_n\}$ are sets in I so $\{n \in \mathbb{N}: \Delta^m x_k \neq \Delta^m y_k \text{ for some } k \in I_n\} \cup A(\varepsilon, \mu) \in I$, which immediately gives

$$\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m y_k - \mathcal{L}, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m y_k - \mathcal{L}, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m y_k - \mathcal{L}, \mu) \geq \varepsilon \right\} \in I ;$$

and therefore (y_k) is $\Delta^m(I_\lambda) - \text{convergent}$. ■.

5. $\Delta^m(I_\lambda) - \text{limit points}$ and $\Delta^m(I_\lambda) - \text{cluster points}$

Definition 5.1 An element \mathcal{L}_0 is said to be a $\Delta^m - \text{limit point}$ of (x_k) w.r.t \mathcal{N} if \exists a subsequence of $(\Delta^m x_k)$ that is convergent to \mathcal{L}_0 w.r.t \mathcal{N} .

Definition 5.2 An element \mathcal{L}_0 is said to be a $\Delta^m(I_\lambda) - \text{limit point}$ of (x_k) if $\exists, K = \{k_1 < k_2 < \dots < k_j < \dots\} \subseteq \mathbb{N}$ s.t the set $K' = \{r \in \mathbb{N}: k_j \in I_r\} \notin I$ and $\mathcal{N}_\lambda - \lim_j \Delta^m x_{k_j} = \mathcal{L}_0$.

Let $\Lambda_{\mathcal{N}}^\lambda(I, x)$ denotes the set of all $\Delta^m(I_\lambda) - \text{limit points}$ of $x = (x_k)$.

Definition 5.3 An element \mathcal{L}_0 is said to be a $\Delta^m(I_\lambda) - \text{cluster point}$ of (x_k) if $0 < \varepsilon < 1$ and $\mu > 0$,

$$\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_0, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \right\} \notin I.$$

Let $\Gamma_N^\lambda(I, x)$ denotes the set of $\Delta^m(I_\lambda)$ – cluster points of the sequence $x = (x_k)$.

Theorem 5.1 For $x = (x_k)$ in V , $\Lambda_N^\lambda(I, x) \subseteq \Gamma_N^\lambda(I, x)$.

Proof Let $\mathcal{L}_0 \in \Lambda_N^\lambda(I, x)$, \exists set $K = \{k_1 < k_2 < \dots < k_j < \dots\} \subseteq \mathbb{N}$ s.t. $K' = \{r \in \mathbb{N}: k_j \in I_r\} \notin I$, $\mathcal{N}_\lambda - \lim_j \Delta^m x_{k_j} = \mathcal{L}_0$. So for $0 < \varepsilon < 1$ and $\mu > 0$, $\exists, j_0 \in \mathbb{N}$ s.t. for $j \geq j_0$

$$\frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{G}(\Delta^m x_{k_j} - \mathcal{L}_0, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{B}(\Delta^m x_{k_j} - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{Y}(\Delta^m x_{k_j} - \mathcal{L}_0, \mu) < \varepsilon$$

This implies that

$$\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_0, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \right\} \subseteq K' - \{k_1 < k_2 < \dots < k_{j_0}\}$$

Since I is admissible so $K' - \{k_1 < k_2 < \dots < k_{j_0}\} \notin I$, which immediately gives

$$\left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_0, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \right\} \notin I;$$

and therefore $\mathcal{L}_0 \in \Gamma_N^\lambda(I, x)$. ■

Theorem 5.2 For $x = (x_k)$ in V , $\Gamma_N^\lambda(I, x)$ is a closed set..

Proof. Let $\mathcal{L}_0 \in \overline{\Gamma_N^\lambda(I, x)}$ where bar denotes the closure of the set. Let, $0 < \varepsilon < 1$ and $\mu > 0$. Since $\mathcal{L}_0 \in \overline{\Gamma_N^\lambda(I, x)}$ so $\Gamma_N^\lambda(I, x) \cap O(\mathcal{L}_0, \varepsilon, \mu)$ where $O(\mathcal{L}_0, \varepsilon, \mu) = \{\mathcal{L} \in V: \mathcal{G}(\mathcal{L} - \mathcal{L}_0, \mu) > 1 - \varepsilon, \mathcal{B}(\mathcal{L} - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \mathcal{Y}(\mathcal{L} - \mathcal{L}_0, \mu) < \varepsilon\}$ denotes the open neighborhood of \mathcal{L}_0 . Let $\mathcal{L}_1 \in \Gamma_N^\lambda(I, x) \cap O(\mathcal{L}_0, \varepsilon, \mu)$. Choose $\gamma > 0$ such that $O(\mathcal{L}_1, \gamma, \mu) \subseteq O(\mathcal{L}_0, \varepsilon, \mu)$. Now we have

$$A = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_0, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_0, \mu) < \varepsilon \right\} \supseteq$$

$$\left\{ r \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{G}(\Delta^m x_k - \mathcal{L}_1, \mu) > 1 - \gamma, \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{B}(\Delta^m x_k - \mathcal{L}_1, \mu) < \gamma \text{ and } \frac{1}{\lambda_n} \sum_{j \in I_n} \mathcal{Y}(\Delta^m x_k - \mathcal{L}_1, \mu) < \gamma \right\} = B.$$

As $\mathcal{L}_1 \in \Gamma_{\mathcal{N}}^\lambda(I, x)$ so $B \notin I$, and consequently $A \notin I$. Hence $\mathcal{L}_0 \in \Gamma_{\mathcal{N}}^\lambda(I, x)$ and therefore $\Gamma_{\mathcal{N}}^\lambda(I, x)$ is a closed set in V . ■

6. $\Delta^m(\lambda)$ –Cauchy and $\Delta^m(I_\lambda)$ –Cauchy sequences

Definition 6.1 A sequence $x = (x_k)$ in V is said to be $\Delta^m(\lambda)$ –Cauchy w.r.t \mathcal{N} if \exists positive integers p and n_0 s.t.

$$\frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - x_p, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - x_p, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - x_p, \mu) < \varepsilon \text{ for } n \geq n_0$$

Definition 6.2 Let $0 < \varepsilon < 1$ and $\mu > 0$. A sequence $x = (x_k)$ in V is said to be $\Delta^m(I_\lambda)$ –Cauchy w.r.t \mathcal{N} if \exists positive integers p s.t

$$\left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - x_p, \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - x_p, \mu) \geq \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - x_p, \mu) \geq \varepsilon \right\} \in I$$

or equivalently

$$\left\{ n \in \mathbb{N} : \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - x_p, \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - x_p, \mu) < \varepsilon \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - x_p, \mu) < \varepsilon \right\} \in \mathcal{F}(I).$$

Definition 6.3 Let, $0 < \varepsilon < 1$ and $\mu > 0$. A sequence $x = (x_k)$ in V is said to be $\Delta^m(I_\lambda^*)$ –Cauchy w.r.t \mathcal{N} if $\exists, K = \{k_1 < k_2 < \dots < k_j < \dots\} \subseteq \mathbb{N}$ s.t. $K' = \{n \in \mathbb{N} : k_j \in I_n\} \in \mathcal{F}(I)$ and (x_{k_j}) is $\Delta^m(\lambda)$ –Cauchy sequence w.r.t \mathcal{N} .

The following two Theorems are analogs of Theorem 4.3, and Theorem 3.3 respectively.

Theorem 6.1 2 If $x = (x_k)$ in V is $\Delta^m(\lambda)$ –Cauchy then it is $\Delta^m(I_\lambda)$ –Cauchy.

Theorem 6.2 If $x = (x_k)$ in V is $\Delta^m(\lambda)$ –Cauchy then \exists a subsequence of $(\Delta^m x_k)$ that is \mathcal{N} –Cauchy.

Theorem 6.3 If $x = (x_k)$ in V is $\Delta^m(I_\lambda^*)$ –Cauchy, then it is $\Delta^m(I_\lambda)$ –Cauchy.

7. $\Delta^m(I_\lambda)$ – Convergence preserving linear operators

In this section, we define continuous and sequentially continuous mappings and present a characterization of I_λ – convergence preserving linear operators.

Definition 7.1 A mapping $T: V \rightarrow V$ is said to be continuous at $x_0 \in V$ w.r.t \mathcal{N} if for $0 < \varepsilon < 1$ and $\mu > 0, \exists 0 < \varepsilon' < 1$ and $\gamma > 0$ s.t. for all $x \in V$,

$$\mathcal{G}(x - x_0, \gamma) > 1 - \varepsilon', \quad \mathcal{B}(x - x_0, \gamma) < \varepsilon' \quad \text{and} \quad \mathcal{Y}(x - x_0, \gamma) < \varepsilon'$$

imply that

$$\mathcal{G}(T(x) - T(x_0), \mu) > 1 - \varepsilon, \quad \mathcal{B}(T(x) - T(x_0), \mu) < \varepsilon \quad \text{and} \quad \mathcal{Y}(T(x) - T(x_0), \mu) < \varepsilon.$$

If This is continuous at every point of the space V , then we call T continuous on V .

Definition 7.2 A mapping $T: V \rightarrow V$ is said to be sequentially continuous at $x_0 \in V$ w.r.t \mathcal{N} and Δ^m if for any sequence $x = (x_k)$ in V with $\mathcal{N} - \lim_k \Delta^m x_k = x_0$ implies that $\mathcal{N} - \lim_k T(\Delta^m x_k) = T(x_0)$.

Theorem 7.1 Let V be an NNS. A mapping $T: V \rightarrow V$ is continuous w.r.t \mathcal{N} if and only if it is sequentially continuous with respect to the neutrosophic norm \mathcal{N} .

Proof. Omitted.

Definition 7.3 A mapping $T: V \rightarrow V$ is said to preserve $\Delta^m(I_\lambda)$ – convergence in V if $\Delta^m(I_\lambda^N) - \lim_k x_k = x_0$ implies $\Delta^m(I_\lambda^N) - \lim_k T(x_k) = T(x_0)$ for every sequence $x = (x_k)$ in V which is $\Delta^m(I_\lambda)$ – convergent to $x_0 \in V$.

Theorem 7.2. A linear operator $T: V \rightarrow V$ preserves $\Delta^m(I_\lambda)$ – convergence in V if and only if T is continuous on V .

Proof First suppose that T is continuous on V . Let $x = (x_k)$ be any sequence in V such that $\Delta^m(I_\lambda^N) - \lim_k x_k = x_0$. For each $0 < \varepsilon < 1$ and $\mu > 0$, there exists $0 < \varepsilon' < 1$ and $\gamma > 0$ such that for all $x \in V$,

$$\mathcal{G}(x - x_0, \gamma) > 1 - \varepsilon', \quad \mathcal{B}(x - x_0, \gamma) < \varepsilon' \quad \text{and} \quad \mathcal{Y}(x - x_0, \gamma) < \varepsilon'$$

imply that

$$\mathcal{G}(T(x) - T(x_0), \mu) > 1 - \varepsilon, \quad \mathcal{B}(T(x) - T(x_0), \mu) < \varepsilon \quad \text{and} \quad \mathcal{Y}(T(x) - T(x_0), \mu) < \varepsilon.$$

If,

$$O(x_0, \varepsilon', \gamma) = \{x \in V: \mathcal{G}(x - x_0, \gamma) > 1 - \varepsilon', \mathcal{B}(x - x_0, \gamma) < \varepsilon' \quad \text{and} \quad \mathcal{Y}(x - x_0, \mu) < \varepsilon'\}$$

$$O(T(x_0), \varepsilon, \mu) = \{T(x) \in V: \mathcal{G}(T(x) - T(x_0), \mu) > 1 - \varepsilon, \mathcal{B}(T(x) - T(x_0), \mu) < \varepsilon \text{ and } \mathcal{Y}(T(x) - T(x_0), \mu) < \varepsilon.\}$$

denotes open balls centered at x_0 and $T(x_0)$ respectively, then for all $x \in V$, if $x \in O(x_0, \varepsilon', \gamma)$ then $T(x) \in O(T(x_0), \varepsilon, \mu)$. But then we have

$$A(\varepsilon', \gamma) = \left\{ n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - x_0, \gamma) > 1 - \varepsilon', \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - x_0, \gamma) < \varepsilon' \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - x_0, \gamma) < \varepsilon' \right\}$$

$$\subseteq \left\{ \begin{array}{l} n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(T(\Delta^m x_k) - T(x_0), \mu) > 1 - \varepsilon, \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(T(\Delta^m x_k) - T(x_0), \mu) < \varepsilon \text{ and} \\ \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(T(\Delta^m x_k) - T(x_0), \mu) < \varepsilon \end{array} \right\}$$

$$= B(\varepsilon, \mu).$$

Since $A(\varepsilon', \gamma) \in \mathcal{F}(I)$ therefore $B(\varepsilon, \mu) \in \mathcal{F}(I)$. Hence, we have $\Delta^m(I_N^\lambda) - \lim_k T(x_k) = T(x_0)$.

Conversely, suppose that the operator $T: V \rightarrow V$ preserves $\Delta^m(I_\lambda) - \text{convergence}$ in V . We prove that T is continuous. Suppose that T is not continuous at some point x_0 in V . Then there is some $0 < \varepsilon < 1$ and $\mu > 0$ such that, for all $0 < \varepsilon' < 1$ and $\gamma > 0$ we have if $x \in O(x_0, \varepsilon', \gamma)$ then $T(x) \notin O(T(x_0), \varepsilon, \mu)$ where $x \in V$. Now we can have a sequence $x = (x_k)$ such that $\mathcal{N} - \lim_k \Delta^m x_k = x_0$ but $\mathcal{N} - \lim_k T(\Delta^m x_k) \neq T(x_0)$. This gives

$$D(\varepsilon', \gamma) = \left\{ \begin{array}{l} n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(\Delta^m x_k - x_0, \gamma) \\ > 1 - \varepsilon', \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(\Delta^m x_k - x_0, \gamma) < \varepsilon' \text{ and } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(\Delta^m x_k - x_0, \gamma) < \varepsilon' \end{array} \right\}$$

$$\subseteq \left\{ \begin{array}{l} n \in \mathbb{N}: \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{G}(T(\Delta^m x_k) - T(x_0), \mu) \leq 1 - \varepsilon \text{ or } \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{B}(T(\Delta^m x_k) - T(x_0), \mu) \geq \varepsilon \text{ and} \\ \frac{1}{\lambda_n} \sum_{k \in I_n} \mathcal{Y}(T(\Delta^m x_k) - T(x_0), \mu) \geq \varepsilon \end{array} \right\}$$

$$= E(\varepsilon, \mu).$$

Since $D(\varepsilon', \gamma) \in \mathcal{F}(I)$ so $E(\varepsilon, \mu) \in \mathcal{F}(I)$ and therefore we have $\Delta^m(I_N^\lambda) - \lim_k T(x_k) \neq T(x_0)$. ■

Conclusion

This paper used difference operators to define some new summability concepts on neutrosophic normed spaces. The paper introduced the concepts of generalized limit point, cluster point and obtain some relationships among these notions. In addition, the paper defined generalized Cauchy sequences on these spaces and present a characterization of new summability method that preserve linear operators on neutrosophic normed spaces.

Funding: “This research received no external funding”

Conflicts of Interest: “The authors declare no conflict of interest.”

References

- [1] K Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 20(1986), 87–96.
- [2] T. Bera and N.K. Mahapatra, Neutrosophic soft linear spaces, Fuzzy Information and Engineering, 9 (2017), 299–324.
- [3] T. Bera and N.K. Mahapatra, Neutrosophic soft normed linear spaces, Neutrosophic Sets and Systems, 23 (2018), 52–71.
- [4] B. Choudhary, Lacunary I -convergent sequences, Real Analysis Exchange, Summer Symposium, 2009, 56–57.
- [5] J. S. Connor, The statistical and strong p -Cesaro convergence of sequences, Analysis, 8 (1988), 47–63.

- [6] P. Debnath, Lacunary ideal convergence in intuitionistic fuzzy normed linear spaces, *Computers and Mathematics with Applications*, 63 (2012), 708–715.
- [7] K. Demirci, I -limit superior and inferior, *Math. Commun.*, 6(2001), 165 - 172.
- [8] K. Dems, On I -Cauchy sequences, *Real Analysis Exchange*, 30(2004), 123 -128.
- [9] H. Fast, Surla convergence statistique, *Coll. Math.*, 2 (1951), 241–244.
- [10] J.A. Fridy, On statistical convergence, *Analysis*, 5 (1985), 301–313.
- [11] J.A. Fridy, C. Orhan, Lacunary statistical convergence, *Pacific J. Math.*, 160 (1993), 43–51.
- [12] B. Hazarika, V. Kumar and B. Lafuerza-Guillén, Generalized ideal convergence in Intuitionistic fuzzy normed linear spaces, *Filomat*, 27 (5), 811-820.
- [13] Soheyb Milles , Abdelkrim Latrech , Omar Barkat. (2020). Completeness and Compactness in Standard Single Valued Neutrosophic Metric Spaces, *International Journal of Neutrosophic Science*, Vol. 12(2): 96-104
- [14] M. Kirisci and Simsek, Neutrosophic normed spaces and statistical convergence, *The Journal of Analysis*, 28(2020), 1059-1073.
- [15] A. Komisarski, Pointwise I -convergence and I^* -convergence in measure of sequences of functions, *J. Math. Anal. Appl.*, 340 (2008), 770–779.
- [16] P. Kostyrko, T. Salat, W. Wilczynski, I -convergence, *Real Anal. Exchange*, 26(2) (2000/2001), 669–686.
- [17] P. Kostyrko, M. Macaj, T. Salat and, M. Slezziak, I -convergence and extremal I -limit points, *Math. Slovaca*, 4(2005), 443 - 464.
- [18] V. Kumar, On I and I^* -convergence of double sequences, *Math. Commun*, 12 (2007), 171–181,
- [19] V. Kumar and B. Lafuerza-Guillén, On ideal convergence of double sequences in probabilistic normed spaces, *Acta Mathematica Sinica*, 29(2012), 1689-1700.
- [20] LEINDLER, L.: Uber die de la Vallee-Pousinsche Summierbarkeit allgemeiner Orthogonalreihen, *Acta Math. Acad. Sci. Hungar.* 16 (1965), 375-387
- [21] M. Mursaleen, S. A. Mohiuddine, and O.H.H. Edley, On ideal convergence of double sequences in Intuitionistic fuzzy normed spaces, *Comput. Math. Appl.*, 59 (2010), 603-611.
- [22] J. H. Park, Intuitionistic fuzzy metric spaces, *Chaos, Solitons & Fractals*, 22 (2004), 1039–46.
- [23] R Saadati, J H Park. On the Intuitionistic fuzzy topological spaces, *Chaos, Solitons & Fractals*, 27(2006), 331–44.
- [24] T. Salat, On statistically convergent sequences of real numbers, *Math. Slovaca*, 30 (1980), 139–150.
- [25] I. J. Schoenberg, The inerrability of certain functions and related summability methods, *Amer. Math. Monthly*, 66 (1959), 361–375.
- [26] B. Schweizer, A. Sklar, Statistical metric spaces. *Pacific J Math* 10(1960), 314–44.
- [27] F. Smarandache, Neutrosophic set, a generalization of the intuitionistic fuzzy sets, *International Journal of Pure and Applied Mathematics*, 24(2005), 287–297.
- [28] L.A. Zadeh Fuzzy sets, *Inform Control*, 8(1965), 338–353.