



Generating Weak Fuzzy Complex and Anti Weak Fuzzy Complex Integer Solutions for Pythagoras Diophantine Equation $X^2 + Y^2 = Z^2$

Abuobida M. A. Alfahal^{1,*}, Mohammad Abobala², Yaser Ahmad Alhasan³, Raja Abdullah Abdulfatah⁴

^{1,3,4}Deanship of the Preparatory Year, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia

²Tishreen University, Department Of Mathematics, Latakia, Syria

Emails: a.alfahal@psau.edu.sa; mohammadabobala777@gmail.com; y.alhasan@psau.edu.sa; r.abdulfatah@psau.edu.sa

*Corresponding author: a.alfahal@psau.edu.sa

Abstract

In this paper, we find necessary and sufficient conditions for a weak fuzzy complex integer triple (X, Y, Z) to be a pythagoras triple, and for an anti-weak fuzzy complex integer triple (X, Y, Z) to be a Pythagoras triple, where we prove that the non-linear Fermat's Diophantine equation has three different types of solutions according to the value of ε^2 . All types will be solved and discussed in terms of theorems and examples that explain how the algorithms work.

Keywords: weak fuzzy complex integer; anti-weak fuzzy complex integer; Pythagoras triple; Pythagoras Diophantine equation.

1. Introduction.

The concept of weak fuzzy numbers was defined in [1] as a new generalization of real numbers in a similar approach of split-complex numbers [2-4]. This generalization has many applications and relationships with other algebraic structures. Pythagoras triples are solutions of the non-linear Diophantine equation $X^2 + Y^2 = Z^2$, they were studied by many authors with many interesting results and generalizations with many applications in algebra and computer science, see [5, 8-12].

Weak fuzzy complex numbers were very useful in the study of other algebraic structures such as spaces and products [6-7].

In this paper, we close a research gap by solving the non-linear Diophantine equation $X^2 + Y^2 = Z^2$ in the set of integer weak fuzzy complex numbers, where we discuss the all three type of the equation $X^2 + Y^2 = Z^2$ according to the values of $\varepsilon^2 = t \in]0,1[$, and we proved solutions for every possible case. Also, we discuss the anti-weak fuzzy complex integer solutions for the same equation for $\varepsilon^2 = t > 1$.

For definitions and basic concepts of weak fuzzy complex numbers/integers, see [1, 6-7].

2. Main discussion

Definition:

A weak fuzzy complex Pythagoras triple (X, Y, Z) is defined as follows:

$$X^2 + Y^2 = Z^2; X = x_1 + x_2\varepsilon, Y = y_1 + y_2\varepsilon, Z = z_1 + z_2\varepsilon, \text{ where } x_i, y_i, z_i \in \mathbb{Z}, \varepsilon^2 = t \in]0,1[.$$

Theorem.

The weak fuzzy complex Diophantine equation $X^2 + Y^2 = Z^2$ is equivalent to:

$$\begin{cases} (x_1 + \sqrt{t}x_2)^2 + (y_1 + \sqrt{t}y_2)^2 = (z_1 + \sqrt{t}z_2)^2 \\ (x_1 - \sqrt{t}x_2)^2 + (y_1 - \sqrt{t}y_2)^2 = (z_1 - \sqrt{t}z_2)^2 \end{cases}$$

Proof.

We have:

$$X^2 = (x_1 + x_2\varepsilon)^2 = x_1^2 + x_2^2t + 2x_1x_2\varepsilon$$

$$Y^2 = (y_1 + y_2\varepsilon)^2 = y_1^2 + y_2^2t + 2y_1y_2\varepsilon$$

$$Z^2 = (z_1 + z_2\varepsilon)^2 = z_1^2 + z_2^2t + 2z_1z_2\varepsilon$$

Thus,

$$\begin{cases} x_1^2 + x_2^2t + y_1^2 + y_2^2t = z_1^2 + z_2^2t & (1) \\ 2x_1x_2 + 2y_1y_2 = 2z_1z_2 & (2) \end{cases}$$

We multiply (2) by \sqrt{t} , and compute (1) + \sqrt{t} (2), (1) - \sqrt{t} (2), then:

$$\begin{cases} x_1^2 + x_2^2t + 2\sqrt{t}x_1x_2 + y_1^2 + y_2^2t + 2\sqrt{t}y_1y_2 = z_1^2 + z_2^2t + 2\sqrt{t}z_1z_2 \\ x_1^2 + x_2^2t - 2\sqrt{t}x_1x_2 + y_1^2 + y_2^2t - 2\sqrt{t}y_1y_2 = z_1^2 + z_2^2t - 2\sqrt{t}z_1z_2 \end{cases}$$

Which is equivalent to:

$$\begin{cases} (x_1 + \sqrt{t}x_2)^2 + (y_1 + \sqrt{t}y_2)^2 = (z_1 + \sqrt{t}z_2)^2 \\ (x_1 - \sqrt{t}x_2)^2 + (y_1 - \sqrt{t}y_2)^2 = (z_1 - \sqrt{t}z_2)^2 \end{cases}$$

Thus, the proof is complete.

Definition.

A weak fuzzy complex Pythagoras triple (X, Y, Z) is called of type (1) if $\sqrt{t} = \frac{a}{b} \in Q, gcd(a, b) = 1$.

It is called of type (2) if $\sqrt{t} \notin Q; t = \frac{a}{b} \in Q, gcd(a, b) = 1$.

It is called of type (3) if $\sqrt{t} \notin Q; t \notin Q$.

3. Algorithm for finding solutions for type (1) equation.

Let $\sqrt{t} = \frac{a}{b} \in Q, gcd(a, b) = 1$, then $X^2 + Y^2 = Z^2$ is equivalent to:

$$\begin{cases} \left(x_1 + \frac{a}{b}x_2\right)^2 + \left(y_1 + \frac{a}{b}y_2\right)^2 = \left(z_1 + \frac{a}{b}z_2\right)^2 & (1) \\ \left(x_1 - \frac{a}{b}x_2\right)^2 + \left(y_1 - \frac{a}{b}y_2\right)^2 = \left(z_1 - \frac{a}{b}z_2\right)^2 & (2) \end{cases}$$

We multiply (1), (2) by $b^2 \neq 0$, to get:

$$\begin{cases} (bx_1 + ax_2)^2 + (by_1 + ay_2)^2 = (bz_1 + az_2)^2 & (1) \\ (bx_1 - ax_2)^2 + (by_1 - ay_2)^2 = (bz_1 - az_2)^2 & (2) \end{cases}$$

This mean that:

$$T_1 = (bx_1 + ax_2, by_1 + ay_2, bz_1 + az_2)$$

$$T_2 = (bx_1 - ax_2, by_1 - ay_2, bz_1 - az_2)$$

Are two classical Pythagoras triples.

Now, let $T_1 = (X_1, Y_1, Z_1), T_2 = (X_2, Y_2, Z_2)$ are two Pythagoras triples in Z , then they give us a weak fuzzy complex Pythagoras triple if and only if:

$$\begin{cases} 2b \setminus X_1 + X_2, 2a \setminus X_1 - X_2 \\ 2b \setminus Y_1 + Y_2, 2a \setminus Y_1 - Y_2 \\ 2b \setminus Z_1 + Z_2, 2a \setminus Z_1 - Z_2 \end{cases}$$

on the other hand, we get:

$$X = \left(\frac{X_1 + X_2}{2b}\right) + \varepsilon \left(\frac{X_1 - X_2}{2a}\right), Y = \left(\frac{Y_1 + Y_2}{2b}\right) + \varepsilon \left(\frac{Y_1 - Y_2}{2a}\right), Z = \left(\frac{Z_1 + Z_2}{2b}\right) + \varepsilon \left(\frac{Z_1 - Z_2}{2a}\right)$$

example.

Consider $\varepsilon^2 = t = \frac{1}{9}, \sqrt{t} = \frac{1}{3} \in Q$, then $X^2 + Y^2 = Z^2$ is Pythagoras Diophantine equation of type (1).

Consider the following two classical Pythagoras triples.

$T_1 = (6, 8, 10), T_2 = (24, 10, 26)$ are two pythagoras triples in Z , where $X_1 = 6, Y_1 = 8, Z_1 = 10, X_2 = 24, Y_2 = 10, Z_2 = 26$, and:

$$\begin{cases} X_1 + X_2 = 30, X_1 - X_2 = -18 \\ Y_1 + Y_2 = 18, Y_1 - Y_2 = -2 \\ Z_1 + Z_2 = 36, Z_1 - Z_2 = -16 \end{cases}$$

also,

$$\begin{cases} 2b = 6 \setminus 30, 2a = 2 \setminus -18 \\ 2b = 6 \setminus 18, 2a = 2 \setminus -2 \\ 2b = 6 \setminus 36, 2a = 2 \setminus -16 \end{cases}$$

so that (T_1, T_2) gives a weak fuzzy complex pythagoras triple as follows:

$$X = \left(\frac{30}{6}\right) + \varepsilon \left(\frac{-18}{2}\right) = 5 - 9\varepsilon$$

$$Y = \left(\frac{18}{6}\right) + \varepsilon \left(\frac{-2}{2}\right) = 3 - \varepsilon$$

$$Z = \left(\frac{36}{6}\right) + \varepsilon \left(\frac{-16}{2}\right) = 6 - 8\varepsilon$$

4. Algorithm for solving type (3) equation.

Consider $t \notin Q$, then $X^2 + Y^2 = Z^2$ is written as follows:

$$\begin{cases} x_1^2 + y_1^2 + t(x_2^2 + y_2^2) = z_1^2 + tz_2^2 & (1) \\ x_1x_2 + y_1y_2 = z_1z_2 & (2) \end{cases}$$

Since $x_i, y_i, z_i \in Z$ and $t \in R - Q$, then:

$$\begin{cases} x_1^2 + y_1^2 = z_1^2 & (1) \\ x_2^2 + y_2^2 = z_2^2 & (2) \\ x_3^2 + y_3^2 = z_3^2 & (3) \end{cases}$$

this means that $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are two classical Pythagoras triples in Z .

From equation (3), we get:

$(x_1x_2 + y_1y_2)^2 = z_1^2z_2^2$, hence
 $x_1^2x_2^2 + y_1^2y_2^2 + 2x_1x_2y_1y_2 = z_1^2z_2^2$, thus:
 $x_1^2x_2^2 + (z_1^2 - x_1^2)(z_2^2 - x_2^2) + 2x_1x_2y_1y_2 = z_1^2z_2^2$,

So that:

$$\begin{aligned} 2x_1^2x_2^2 - z_1^2x_2^2 - x_1^2z_2^2 + 2x_1x_2y_1y_2 &= 0, \\ 2x_1x_2(x_1x_2 + y_1y_2) - x_1^2z_2^2 - z_1^2x_2^2 &= 0, \\ 2x_1x_2(z_1z_2) - x_1^2z_2^2 - z_1^2x_2^2 &= 0 \\ (x_1z_2 - x_2z_1)^2 &= 0 \implies \frac{x_1}{x_2} = \frac{z_1}{z_2} \end{aligned}$$

On the other hand, we have:

$x_1^2z_2^2 = z_1^2x_2^2 \implies x_1^2(x_2^2 + y_2^2) = x_2^2(x_1^2 + y_1^2)$, thus:
 $x_1^2y_2^2 = y_1^2x_2^2$, hence $\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2}$ under the assumption of $x_1, x_2, y_1, y_2, z_1, z_2 > 0$.

From this point of view, if $T_1 = (X_1, Y_1, Z_1), T_2 = (X_2, Y_2, Z_2)$ are two classical Pythagoras triples in Z , then they give us a weak fuzzy complex Pythagoras triple of type (3) if and only if:

$$\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = c \in Q.$$

Example.

For $t = \frac{1}{\sqrt{3}} \notin Q$, then $X^2 + Y^2 = Z^2$ is an equation of type (3).

Consider the following Pythagoras triples in $Z, T_1 = (3,4,5), T_2 = (6,8,10)$, then $\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = \frac{1}{2} \in Q$.

This implies that (T_1, T_2) give us a weak fuzzy complex Pythagoras triple with respect to $t = \frac{1}{3}$.

$$X = 3 + 6\varepsilon, Y = 4 + 8\varepsilon, Z = 5 + 10\varepsilon.$$

5. Algorithm for solving equation of type (2).

Consider that $\frac{a}{b} = t \in Q, \sqrt{t} \notin Q$, then $X^2 + Y^2 = Z^2$ is of type (2), and it is written as follows:

$$\begin{cases} x_1^2 + y_1^2 + t(x_2^2 + y_2^2) = z_1^2 + tz_2^2 \\ x_1x_2 + y_1y_2 = z_1z_2 \end{cases}$$

Which equivalent to:

$$\begin{cases} b(x_1^2 + y_1^2 - z_1^2) + a(x_2^2 + y_2^2 - z_2^2) = 0 & (1) \\ x_1x_2 + y_1y_2 = z_1z_2 & (2) \end{cases}$$

Put $L_1 = x_1^2 + y_1^2 - z_1^2, L_2 = x_2^2 + y_2^2 - z_2^2$, then (1) takes the following formula:

$$bL_1 + aL_2 = 0; gcd(a, b) = 1.$$

It is a linear Diophantine equation in two variables L_1, L_2 .

It is solvable since $gcd(a, b) = 1$, and $L_1 = ka, L_2 = -kb$, where $k \in Z$, hence:

$$\frac{L_1}{L_2} = -\frac{a}{b} = -t \text{ if } L_1, L_2 \neq 0, \text{ or } L_1 = L_2 = 0.$$

We have possible cases:

Case (1).

If $L_1 = L_2 = 0$, then $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are two Pythagoras triple in Z with $\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = c \in Q$ (the proof is exactly similar of type (3) equation).

Case (2).

If $L_1 \neq 0, L_2 \neq 0$, then:

$$\begin{cases} L_1 = -\frac{a}{b} = -t < 0 \end{cases} \quad (1)$$

$$\begin{cases} x_1x_2 + y_1y_2 = z_1z_2 \end{cases} \quad (2)$$

Then, $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are not Pythagoras triples in Z .

Example.

For $t = \frac{1}{10}, \sqrt{t} \notin Q$, consider the following triples:

$$T_1 = (X_1, Y_1, Z_1) = (1, 3, 2), T_2 = (X_2, Y_2, Z_2) = (2, 6, 10)$$

We have:

$$\frac{L_1}{L_2} = \frac{x_1^2 + y_1^2 - z_1^2}{x_2^2 + y_2^2 - z_2^2} = \frac{6}{-60} = \frac{-1}{10} = -t$$

Also, $x_1x_2 + y_1y_2 = 20 = z_1z_2$, this implies that:

$X = 1 + 2\varepsilon, Y = 3 + 6\varepsilon, Z = 2 + 10\varepsilon$ is a Pythagoras weak fuzzy complex triple of type (2).

By easy computing, we find:

$$X^2 = 1 + \frac{4}{10} + 4\varepsilon = \frac{14}{10} + 4\varepsilon$$

$$Y^2 = 9 + \frac{36}{10} + 36\varepsilon = \frac{126}{10} + 36\varepsilon$$

$$Z^2 = 4 + \frac{100}{10} + 40\varepsilon = 14 + 40\varepsilon$$

It is easy to see that $X^2 + Y^2 = Z^2$.

Definition:

An anti-weak fuzzy complex Pythagoras triple (X, Y, Z) is defined as follows:

$$X^2 + Y^2 = Z^2; X = x_1 + x_2\varepsilon, Y = y_1 + y_2\varepsilon, Z = z_1 + z_2\varepsilon, \text{ where } x_i, y_i, z_i \in Z, \varepsilon^2 = t > 1.$$

Theorem.

The anti-weak fuzzy complex Diophantine equation $X^2 + Y^2 = Z^2$ is equivalent to:

$$\begin{cases} (x_1 + \sqrt{t}x_2)^2 + (y_1 + \sqrt{t}y_2)^2 = (z_1 + \sqrt{t}z_2)^2 \\ (x_1 - \sqrt{t}x_2)^2 + (y_1 - \sqrt{t}y_2)^2 = (z_1 - \sqrt{t}z_2)^2 \end{cases}$$

Proof.

We have:

$$X^2 = (x_1 + x_2\varepsilon)^2 = x_1^2 + x_2^2t + 2x_1x_2\varepsilon$$

$$Y^2 = (y_1 + y_2\varepsilon)^2 = y_1^2 + y_2^2t + 2y_1y_2\varepsilon$$

$$Z^2 = (z_1 + z_2\varepsilon)^2 = z_1^2 + z_2^2t + 2z_1z_2\varepsilon$$

Thus,

$$\begin{cases} x_1^2 + x_2^2t + y_1^2 + y_2^2t = z_1^2 + z_2^2t \end{cases} \quad (1)$$

$$\begin{cases} 2x_1x_2 + 2y_1y_2 = 2z_1z_2 \end{cases} \quad (2)$$

We multiply (2) by \sqrt{t} , and compute $(1) + \sqrt{t}(2), (1) - \sqrt{t}(2)$, then:

$$\begin{cases} x_1^2 + x_2^2t + 2\sqrt{t}x_1x_2 + y_1^2 + y_2^2t + 2\sqrt{t}y_1y_2 = z_1^2 + z_2^2t + 2\sqrt{t}z_1z_2 \\ x_1^2 + x_2^2t - 2\sqrt{t}x_1x_2 + y_1^2 + y_2^2t - 2\sqrt{t}y_1y_2 = z_1^2 + z_2^2t - 2\sqrt{t}z_1z_2 \end{cases}$$

Which is equivalent to:

$$\begin{cases} (x_1 + \sqrt{t}x_2)^2 + (y_1 + \sqrt{t}y_2)^2 = (z_1 + \sqrt{t}z_2)^2 \\ (x_1 - \sqrt{t}x_2)^2 + (y_1 - \sqrt{t}y_2)^2 = (z_1 - \sqrt{t}z_2)^2 \end{cases}$$

Thus, the proof is complete.

Definition.

An anti-weak fuzzy complex Pythagoras triple (X, Y, Z) is called of type (1) if $\sqrt{t} = \frac{a}{b} \in Q, gcd(a, b) = 1$.

It is called of type (2) if $\sqrt{t} \notin Q; t = \frac{a}{b} \in Q, gcd(a, b) = 1$.

It is called of type (3) if $\sqrt{t} \notin Q; t \notin Q$.

6. Algorithm for finding solutions for type (1) equation.

Let $\sqrt{t} = \frac{a}{b} \in Q, gcd(a, b) = 1$, then $X^2 + Y^2 = Z^2$ is equivalent to:

$$\begin{cases} \left(x_1 + \frac{a}{b}x_2\right)^2 + \left(y_1 + \frac{a}{b}y_2\right)^2 = \left(z_1 + \frac{a}{b}z_2\right)^2 \end{cases} \quad (1)$$

$$\begin{cases} \left(x_1 - \frac{a}{b}x_2\right)^2 + \left(y_1 - \frac{a}{b}y_2\right)^2 = \left(z_1 - \frac{a}{b}z_2\right)^2 \end{cases} \quad (2)$$

We multiply (1), (2) by $b^2 \neq 0$, to get:

$$(bx_1 + ax_2)^2 + (by_1 + ay_2)^2 = (bz_1 + az_2)^2 \quad (1)$$

$$(bx_1 - ax_2)^2 + (by_1 - ay_2)^2 = (bz_1 - az_2)^2 \quad (2)$$

This means that:

$$T_1 = (bx_1 + ax_2, by_1 + ay_2, bz_1 + az_2)$$

$$T_2 = (bx_1 - ax_2, by_1 - ay_2, bz_1 - az_2)$$

Are two classical Pythagoras triples.

Now, let $T_1 = (X_1, Y_1, Z_1), T_2 = (X_2, Y_2, Z_2)$ are two Pythagoras triples in Z , then they give us an anti-weak fuzzy complex Pythagoras triple if and only if:

$$\begin{cases} 2b \setminus X_1 + X_2, 2a \setminus X_1 - X_2 \\ 2b \setminus Y_1 + Y_2, 2a \setminus Y_1 - Y_2 \\ 2b \setminus Z_1 + Z_2, 2a \setminus Z_1 - Z_2 \end{cases}$$

on the other hand, we get:

$$X = \left(\frac{X_1 + X_2}{2b}\right) + \varepsilon \left(\frac{X_1 - X_2}{2a}\right), Y = \left(\frac{Y_1 + Y_2}{2b}\right) + \varepsilon \left(\frac{Y_1 - Y_2}{2a}\right), Z = \left(\frac{Z_1 + Z_2}{2b}\right) + \varepsilon \left(\frac{Z_1 - Z_2}{2a}\right)$$

Example.

Consider $\varepsilon^2 = t = 9, \sqrt{t} = 2 \in Q$, then $X^2 + Y^2 = Z^2$ is Pythagoras Diophantine equation of type (1).

Consider the following two classical Pythagoras triples.

$T_1 = (6,8,10), T_2 = (10,24,26)$ are two pythagoras triples in Z , where $X_1 = 6, Y_1 = 8, Z_1 = 10, X_2 = 24, Y_2 = 10, Z_2 = 26$, and:

$$\begin{cases} X_1 + X_2 = 16, X_1 - X_2 = -4 \\ Y_1 + Y_2 = 32, Y_1 - Y_2 = -16 \\ Z_1 + Z_2 = 36, Z_1 - Z_2 = -16 \end{cases}$$

also,

$$\begin{cases} 2b = 2 \setminus 16, 2a = 4 \setminus -4 \\ 2b = 2 \setminus 32, 2a = 4 \setminus -16 \\ 2b = 2 \setminus 36, 2a = 4 \setminus -16 \end{cases}$$

so that (T_1, T_2) gives an anti-weak fuzzy complex pythagoras triple as follows:

$$X = \left(\frac{16}{2}\right) + \varepsilon \left(\frac{-4}{4}\right) = 8 - \varepsilon$$

$$Y = \left(\frac{32}{2}\right) + \varepsilon \left(\frac{-16}{4}\right) = 16 - 4\varepsilon$$

$$Z = \left(\frac{36}{2}\right) + \varepsilon \left(\frac{-16}{4}\right) = 18 - 4\varepsilon$$

7. Algorithm for solving type (3) equation.

Consider $t \notin Q$, then $X^2 + Y^2 = Z^2$ is written as follows:

$$\begin{cases} x_1^2 + y_1^2 + t(x_2^2 + y_2^2) = z_1^2 + tz_2^2 \quad (1) \\ x_1x_2 + y_1y_2 = z_1z_2 \quad (2) \end{cases}$$

Since $x_i, y_i, z_i \in Z$ and $t \in R - Q$, then:

$$\begin{cases} x_1^2 + y_1^2 = z_1^2 \quad (1) \\ x_2^2 + y_2^2 = z_2^2 \quad (2) \\ x_3^2 + y_3^2 = z_3^2 \quad (3) \end{cases}$$

This means that $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are two classical Pythagoras triples in Z .

From equation (3), we get:

$$(x_1x_2 + y_1y_2)^2 = z_1^2z_2^2, \text{ hence}$$

$$x_1^2x_2^2 + y_1^2y_2^2 + 2x_1x_2y_1y_2 = z_1^2z_2^2, \text{ thus:}$$

$$x_1^2x_2^2 + (z_1^2 - x_1^2)(z_2^2 - x_2^2) + 2x_1x_2y_1y_2 = z_1^2z_2^2,$$

So that:

$$2x_1^2x_2^2 - z_1^2x_2^2 - x_1^2z_2^2 + 2x_1x_2y_1y_2 = 0,$$

$$2x_1x_2(x_1x_2 + y_1y_2) - x_1^2z_2^2 - z_1^2x_2^2 = 0,$$

$$2x_1x_2(z_1z_2) - x_1^2z_2^2 - z_1^2x_2^2 = 0$$

$$(x_1z_2 - x_2z_1)^2 = 0 \implies \frac{x_1}{x_2} = \frac{z_1}{z_2}$$

On the other hand, we have:

$$x_1^2z_2^2 = z_1^2x_2^2 \implies x_1^2(x_2^2 + y_2^2) = x_2^2(x_1^2 + y_1^2), \text{ thus:}$$

$$x_1^2y_2^2 = y_1^2x_2^2, \text{ hence } \frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} \text{ under the assumption of } x_1, x_2, y_1, y_2, z_1, z_2 > 0.$$

From this point of view, if $T_1 = (X_1, Y_1, Z_1), T_2 = (X_2, Y_2, Z_2)$ are two classical Pythagoras triples in Z , then they give us an anti-weak fuzzy complex Pythagoras triple of type (3) if and only if:

$$\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = c \in Q.$$

Example.

For $t = \sqrt{3} \notin Q$, then $X^2 + Y^2 = Z^2$ is an equation of type (3).

Consider the following Pythagoras triples in Z , $T_1 = (3,4,5)$, $T_2 = (6,8,10)$, then $\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = \frac{1}{2} \in Q$.

This implies that (T_1, T_2) give us a weak fuzzy complex Pythagoras triple with respect to $t = 3$.

$$X = 3 + 6\varepsilon, Y = 4 + 8\varepsilon, Z = 5 + 10\varepsilon.$$

8. Algorithm for solving equation of type (2).

Consider that $\frac{a}{b} = t \in Q$, $\sqrt{t} \notin Q$, then $X^2 + Y^2 = Z^2$ is of type (2), and it is written as follows:

$$\begin{cases} x_1^2 + y_1^2 + t(x_2^2 + y_2^2) = z_1^2 + tz_2^2 \\ x_1x_2 + y_1y_2 = z_1z_2 \end{cases}$$

Which equivalent to:

$$b(x_1^2 + y_1^2 - z_1^2) + a(x_2^2 + y_2^2 - z_2^2) = 0 \quad (1)$$

$$\begin{cases} b(x_1^2 + y_1^2 - z_1^2) + a(x_2^2 + y_2^2 - z_2^2) = 0 \\ x_1x_2 + y_1y_2 = z_1z_2 \end{cases} \quad (2)$$

Put $L_1 = x_1^2 + y_1^2 - z_1^2$, $L_2 = x_2^2 + y_2^2 - z_2^2$, then (1) takes the following formula:

$$bL_1 + aL_2 = 0; \gcd(a, b) = 1.$$

It is a linear Diophantine equation in two variables L_1, L_2 .

It is solvable since $\gcd(a, b) = 1$, and $L_1 = ka, L_2 = -kb$, where $k \in Z$, hence:

$$\frac{L_1}{L_2} = -\frac{a}{b} = -t \text{ if } L_1, L_2 \neq 0, \text{ or } L_1 = L_2 = 0.$$

We have possible cases:

Case (1).

If $L_1 = L_2 = 0$, then $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are two Pythagoras triple in Z with $\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2} = c \in Q$ (the proof is exactly similar of type (3) equation).

Case (2).

If $L_1 \neq 0, L_2 \neq 0$, then:

$$\begin{cases} \frac{L_1}{L_2} = -\frac{a}{b} = -t < 0 \\ x_1x_2 + y_1y_2 = z_1z_2 \end{cases} \quad (2)$$

Then, $(x_1, y_1, z_1), (x_2, y_2, z_2)$ are not Pythagoras triples in Z .

Example.

For $t = 10$, $\sqrt{t} \notin Q$, consider the following triples:

$$T_2 = (x_2, y_2, z_2) = (1,3,2), T_1 = (x_1, y_1, z_1) = (2,6,10)$$

We have:

$$\frac{L_1}{L_2} = \frac{x_1^2 + y_1^2 - z_1^2}{x_2^2 + y_2^2 - z_2^2} = \frac{-60}{6} = -10 = -t$$

Also, $x_1x_2 + y_1y_2 = 20 = z_1z_2$, this implies that:

$$X = 2 + \varepsilon, Y = 6 + 3\varepsilon, Z = 10 + 2\varepsilon \text{ is a Pythagoras anti-weak fuzzy complex triple of type (2).}$$

9. Conclusion

In this paper, we studied for the first time the weak fuzzy complex and anti-weak fuzzy complex Pythagoras triples, where algorithms for generating weak fuzzy and anti-weak fuzzy complex integer solutions for the non-linear Diophantine equation $X^2 + Y^2 = Z^2$ are presented and handled by many related theorems and examples that clarify the novelty of this work.

Acknowledgments " This study is supported via funding from Prince sattam bin Abdulaziz University project number (PSAU/2023/R/1445)".

References

- [1] Hatip, A., "An Introduction To Weak Fuzzy Complex Numbers ", Galoitica Journal Of Mathematical Structures and Applications, Vol.3, 2023.
- [2] Ahmad, K., " On Some Split-Complex Diophantine Equations", Neoma Journal Of Mathematics and Computer Science, 2023.
- [3] Khaldi, A., " A Study On Split-Complex Vector Spaces", Neoma Journal Of Mathematics and Computer Science, 2023.

- [4] Merkepci, M., and Abobala, M., " On Some Novel Results About Split-Complex Numbers, The Diagonalization Problem And Applications To Public Key Asymmetric Cryptography", *Journal of Mathematics*, Hindawi, 2023.
- [5] Merkepci, M., and Abobala, M., " Security Model for Encrypting Uncertain Rational Data Units Based on Refined Neutrosophic Integers Fusion and El Gamal Algorithm ", *Fusion: Practice and Applications*, 2023.
- [6] Ali, R., " On The Weak Fuzzy Complex Inner Products On Weak Fuzzy Complex Vector Spaces", *Neoma Journal Of Mathematics and Computer Science*, 2023.
- [7] Hatip, A., "On The Weak Fuzzy Complex Vector Spaces", *Galoitica Journal Of Mathematical Structures And Applications*, Vol.3, 2023.
- [8] Merkepci, M., Abobala, M., and Allouf, A., " The Applications of Fusion Neutrosophic Number Theory in Public Key Cryptography and the Improvement of RSA Algorithm ", *Fusion: Practice and Applications*, 2023.
- [9] Abobala, M., and Allouf, A., " On A Novel Security Scheme for The Encryption and Decryption Of 2×2 Fuzzy Matrices with Rational Entries Based on The Algebra of Neutrosophic Integers and El-Gamal Crypto-System", *Neutrosophic Sets and Systems*, vol.54, 2023.
- [10] Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Concept Of Symbolic 7-Plithogenic Real Matrices", *Pure Mathematics For Theoretical Computer Science*, Vol.1, 2023.
- [11] Ibrahim, M., and Abobala, M., "An Introduction To Refined Neutrosophic Number Theory", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [12] Sankari, H., and Abobala, M., "Neutrosophic Linear Diophantine Equations With two Variables", *Neutrosophic Sets and Systems*, Vol. 38, pp. 22-30, 2020.